

**Length-weight relationships of mesopelagic fishes from the  
equatorial and tropical Atlantic waters: influence of environment  
and body shape**

Cristina López-Pérez<sup>1</sup>, M. Pilar Olivar<sup>1</sup>, P. Alexander Hulley<sup>2,3</sup>, Víctor M.  
Tuset<sup>1</sup>

<sup>1</sup>Instituto de Ciencias del Mar (CSIC), Passeig Marítim 37-49, Barcelona 08003, Spain,

<sup>2</sup>Iziko – South African Museum, Cape Town, South Africa

<sup>3</sup>MA-RE Institute, University of Cape Town, South Africa

Correspondence:

Víctor Manuel Tuset Andújar, Instituto de Ciencias del Mar (CSIC), Passeig Marítim 37-49,  
Barcelona 08003, Spain.

Email: vtuset@icm.csic.es

## Abstract

Length-weight relationships (LWR) were estimated for 36 mesopelagic fish species collected from the equatorial and tropical Atlantic encompassing several oceanographic regions: oligotrophic, equatorial, Cape Blanc, Cape Verde and the Canary Islands. The sample was composed of myctophids (25 specimens), gonostomatids (5), sternoptychids (3), stomiids (2) and phosichthyids (1). The species were clustered according to body shape: ‘short and deep’ (sternoptychids), ‘elongate’ (gonostomids, stomiids and some phosichthyids) and ‘fusiform’ (myctophids and some phosichthyids). Three types of weight and LWRs were considered: wet weight ( $WW$ ), eviscerated wet weight ( $eWW$ ) and eviscerated dry weight ( $eDW$ ). The study demonstrated that most species present a positive allometric growth, independent of the weight used. However, the allometric value varied in 40-50% of species depending on the type of weight considered. Significant variations linked to fish morphology were found in the relationship between the slope and intercept of the LWR equation. Significant differences were also noted in the water content linked to fish body shape. Based on the distributions of several species we compare their fitness between oceanographic regions using the relative condition factor ( $K_{rel}$ ). Except for *Diaphus brachycephalus* (oligotrophic vs equatorial waters) and *Lampanyctus alatus* (equatorial, Cape Blanc, Cape Verde and the Canary Islands), no regional significant differences were observed in the species analysed.

## KEYWORDS

mesopelagic fishes, oceanographic gradient, morphology, size-weight growth, Atlantic Ocean

## 1. INTRODUCTION

Mesopelagic fishes have a worldwide distribution from the Arctic to the Antarctic (Krefft, 1974; Hulley, 1981), although species richness and annual production are commonly greater in subtropical and tropical regions (Gjøsaeter & Kawaguchi, 1980). They are generally small to medium-size fishes, including a high diversity of species, numerically dominated by bristlemouths (family Gonostomatidae) and other Stomiiformes, and by lanternfishes (Order Myctophiformes), the majority of which live in the ocean's twilight zone (by definition between 200 and 1000 m). Some, particularly myctophids, undertake diel vertical migrations following their prey into the epipelagic zone to feed at night (Sutton *et al.*, 2008, 2013; Davison *et al.*, 2013; Bernal *et al.*, 2015; Choy *et al.*, 2015; Wang *et al.*, 2019). During daytime, and mainly to avoid predators, they descend into deeper waters mainly to avoid predators where they undertake digestion and excretion. This results in a substantial contribution to total vertical carbon flux from the surface to the deep ocean (Ikeda *et al.*, 2008; Davison *et al.*, 2013; Hudson *et al.*, 2014; Drazicek and Sutton, 2017). Thus, these fishes play a key role in food web interactions linking primary consumers (e.g., copepods and macro-zooplankton) with both pelagic and deep-sea fish species, as well as other marine species such as marine mammals and birds (Springer *et al.*, 1999; Pereira *et al.*, 2011; Smith *et al.*, 2011; Trueman *et al.*, 2014). The high species richness of stomiiform and myctophiform fishes has contributed to the lack of more detailed information on the basic aspects of fish biology, including length-weight relationships (LWR) and condition factors at the species level (Fock & Ehrich, 2010; Battaglia *et al.*, 2010, 2015; Jiang *et al.*, 2017; Fock & Czudaj, 2018; Wang *et al.*, 2018).

LWR's have been used extensively to gather information on growth, ontogenetic changes, population dynamics and in trophic ecology studies to reconstruct and estimate the biomass of partially digested prey (Pauly, 1993; Verdiell-Cubedo *et al.*, 2006; Battaglia *et al.*, 2010, 2016).

In general, the parameters for LWRs differ among species due to multiple factors, including the fish body shape (Schneider *et al.*, 2000; Kulbicki *et al.*, 2005; Froese, 2006; Jellyman *et al.*, 2013), whilst intra-specific differences may be due to gonad maturity, gender, diet, stomach fullness, health, preservation techniques, season and habitat (Beyer, 1987; Sakuma *et al.*, 1999). Even, the sampling protocols (e.g., fresh, frozen or preserved in solution) can lead to variations in LWR estimations (Eduardo *et al.*, 2019). Some applications of LWRs include the ecosystem-modelling approach to obtain confident production-over-biomass estimates (Christensen & Pauly, 1992; Pauly *et al.*, 2000; Christensen & Walters, 2004; Torres *et al.*, 2012). In this sense, LWR's from similar species and different regions are commonly used for the biomass estimation of mesopelagic fish species from acoustics (Fock & Ehrich, 2010; Fock & Czudaj, 2018), despite possible differences in allometric relationships (Atkinson, 1989). Similarly, the condition factor is considered as a proxy of fitness in determining the population health and variability of individuals and populations (Millar & Hickling, 1990; Lloret *et al.*, 2002, 2014; Pazianoto *et al.*, 2016; Brosset *et al.*, 2018). But there is a lack of information for mesopelagic fishes (Watanabe & Kawaguchi, 2003).

The subtropical western Atlantic Ocean is one of the most oligotrophic regions of the global ocean (Morel *et al.*, 2010), whereas the eastern central Atlantic is one of the four most productive regions of the world, due to upwelling (Mittelstaedt, 1983). The ocean between these two regions is characterized by the convergence of water masses originating in both the southern and northern hemispheres, and results in a complex oceanic current system (Stramma & Schott, 1999). Moreover, south of the Cape Verde archipelago, the mesopelagic layers are characterized by the presence of an oxygen minimum zone (OMZ), with dissolved oxygen concentrations corresponding to intermediate hypoxia values (Ekau *et al.*, 2010; Moffitt *et al.*, 2014; Olivar *et al.*, 2017). Previous investigations in the area have shown that some mesopelagic fishes are associated with particular waters masses, while others occur across the

entire region (Olivar *et al.*, 2017, 2018). These differences in the oceanographic features may also have an influence on the trophic transfer efficiencies from plankton to fishes, which can be indirectly evaluated from body condition (Le Cren, 1951; Simpkins & Hubert, 2000; Stevenson & Woods 2006; Wilson & Nussey, 2010).

This study attempts: (i) to provide information on LWR's of 36 mesopelagic fish species collected in the tropical and equatorial Atlantic waters by using different weight variables, so as to facilitate effective use by scientists working on mesopelagic biomass estimation; (ii) to assess the changes in the fish health between oceanographic regions using the Relative Condition Factor (Le Cren, 1951); and (iii) to evaluate how the body shape influences the  $a$  and  $b$  parameters of LWR equation, as well as the water content of species. These types of information can provide useful evidence to improve the understanding of how species and individuals are structured in open waters.

## 2. MATERIAL AND METHODS

### 2.1. Sample collection

Specimens were collected during a cruise carried out in April 2015 across the equatorial and tropical Atlantic, from near the Brazilian coast (13°S 38.3°W) to south Canary Islands (28°N 15.6°E), on board R/V Hespérides. A total of 12 stations were occupied both day and night, and fish samples were obtained in different strata of the water column, from 800 m to the surface (Olivar *et al.*, 2017). Ship speed was 2 knots (1 m/s) and haul duration at each strata ranged from 10 to 30 min. The sampling gear deployed was a pelagic midwater trawl, the “Mesopelagos” net (35-m<sup>2</sup> mouth opening; total length 58 m, with graded-mesh netting from 30 mm at mouth to 4 mm towards the cod end) (more details in Olivar *et al.*, 2017).

The fresh samples were analysed on board. Fishes were identified to the lowest possible taxon and kept frozen at -20°C until transference to the laboratory, where their identifications were checked, and individual measurements of length and weight were taken. Standard length (*SL*) was recorded to the nearest 1 mm, using digital calipers, and whole body wet weight (*WW*) to the nearest 0.0001 g, using a digital balance. Individuals were then dissected, the digestive tract, stomach and gonads were removed, and the eviscerated wet weight (*eWW*) of each specimen was taken. To eliminate water content, eviscerated specimens were freeze-dried in a Telstar LyoAlfa 6 freeze dryer for 24 h, and the eviscerated dry weight of each specimen was then recorded (*eDW*). Water content was calculated as the difference between *eWW* and *eDW*, expressed as % *eWW*, to highlight their effect on the estimation of LWRs.

## 2.2. Length-weight relationships

Size and weight data were fitted using a power function:  $y = ax^b$ , where  $x$  is *SL* and  $y$  the weight parameter (*WW*, *eWW* or *eDW*). The slope,  $b$ , is the allometric growth factor and the intercept,  $a$ , is the expected value of  $y$  at  $x = 1$  (Gould, 1966). Data from the logarithmically transformed equations were then adjusted by the method of least squares. The values of  $b$ , their 95% confidence intervals (95% CI), and the coefficient of determination ( $r^2$ ) were calculated according to the methods of Sokal & Rohlf (1979). In addition, a Student's- $t$ -test was used to evaluate the isometric growth ( $b = 3$ ), and whether  $b$  value is significantly higher or lower indicating a positive or negative allometry, respectively (Gould, 1966; Margalef, 1974; Pauly, 1984).

To assess the influence of body morphology on the intercept ( $a$ ) and slope ( $b$ ) parameters of the LWR (using *eDW*), a linear regression was estimated for three groups of fishes defined on the basis of their general body forms: 'short and deep' (Sternophthyidae), 'elongate'

(Gonostomatidae and Stomiidae) and ‘fusiform’ (Phosichthyidae and Myctophiidae). Statistical differences between the slopes were tested using an ANCOVA analysis. In addition, the average water content of each fish group was compared using a Kruskal-Wallis test (non-parametric test).

### 2.3. Relative condition factor

The fish condition was estimated using the Relative Condition Factor ( $K_{rel} = (W_o/aSL^b$ ; [Le Cren, 1951](#)), which relates the observed body weight ( $W_o = eDW$ ) of a given individual with the predicted by the length–weight relationship ( $aSL^b$ ). The parameters  $a$  and  $b$  were obtained from the regional LWR derived by pooling data for all regions for each species separately ([Efitre et al., 2009](#)). Calculation of Fulton’s Condition Index ( $Kn = 100W/L^3$ ; [Fulton, 1911](#)) was omitted because the growth of mesopelagic fishes was not isometric (see results). The Relative Weight index ([Wege & Anderson, 1978](#); [Froese, 2006](#)) was not used because there are no studies based on this index supporting a population differentiation of mesopelagic fishes in open ocean.

In order to perform the regional comparisons in the  $K_{rel}$ , regions of the study area were defined according to oceanographic parameters described in [Olivar et al. \(2017\)](#): oligotrophic (stations #1-3), equatorial (stations #4-6), Cape Verde islands (stations #7-10), Cape Blanc upwelling (station#11), Canary Islands (station#12) (Figure 1). For these comparisons, only 9 species (*Chauliodus danae*, *Ceratoscopelus warmingii*, *Diaphus brachycephalus*, *Diaphus mollis*, *Diaphus rafinesquii*, *Lampanyctus alatus*, *Lepidophanes guentheri*, *Myctophum nitidulum* and *Vinciguerrria nimbaria*) were considered to have a sufficient number of individuals ( $n > 10$ ) by region and LWRs with high correlations ( $r^2 > 0.90$ ). The normality and homogeneity of variances in the  $K_{rel}$  data were checked for each species by region using the Shapiro-Wilk’s test and Bartlett’s test, respectively. Depending on the Gaussian distribution of

data, variations of  $K_{rel}$  between two regions were compared by the Student's  $t$ -test (parametric test), or the Mann–Whitney's  $U$  Test (non-parametric test). To analyse more than two regions, a Kruskal-Wallis test followed by *a posteriori* Dunn test was performed with the package *dunn.test* v.1.3.5. in R (Dino, 2017).

All statistical analyses and graphical representations were conducted with the software R (R Core Team, 2016).

### 3. RESULTS

A total 1277 individuals belonging to 36 species from 5 families were analysed. The family Myctophidae was represented by 11 genera and 25 species, followed by Gonostomatidae, with 2 genera and 5 species; Sternoptychidae, with 3 genera and 3 species; Stomiidae, with 2 genera and 2 species, and Phosichthyidae, with 1 genus and 1 species (Table 1).

The determination coefficients ( $r^2$ ) of the fitted equations were generally high (mean value of  $0.991 \pm 0.001$ ), ranging from 0.613 for *Cyclothone pseudopallida* (Gonostomatidae) to 0.997 for *Bolinichthys photothorax* (Myctophidae). Excluding the extreme case presented by *C. pseudopallida*, 77% of the relationship have  $r^2 > 0.950$  (Table 2 for *eDW*; Table S1 for *WW* and Table S2 for *eWW*, Supplementary material).

In terms of growth, the LWRs for *WW* revealed that 57.2% of the species showed isometric growth, the 30.5% positive allometry ( $b > 3$ ) and the 11.1% negative allometry ( $b < 3$ ). A similar tendency was also detected when fitting *eWW* (52.7%, 33.3% and 13.8%, respectively). The main differences in growth patterns between the LWRs for *WW* or *eWW* were observed for *Vinciguerria nimbaria* (Phosichthyidae), which presented a higher  $b$  when fitting total weight, and for *Lepidophanes guentheri* (Myctophidae) and *Sternoptyx diaphana* (Sternoptychidae)



with lower  $b$  using eviscerated data. By contrast, LWRs for  $eDW$  showed a slight increment in the proportion of species with positive allometry (47.2% isometric, and 38.8% and 13.8%, positive and negative allometry, respectively). Irrespective of the fitted data, the lowest allometric coefficients were always found for *Cyclothone* spp., together with the myctophid *Diaphus vanhoeffeni*. The highest allometric growth coefficient was always in *Polyipnus polli*, and in myctophids such as *Benthosema glaciale*, *Ceratoscopelus warmingii*, *Diaphus brachycephalus*, *D. holti*, *D. metopoclampus*, *D. rafinesquii*, *Lampanyctus alatus*, *L. pusillus*, *Lepidophanes guentheri*, *Lobianchia dofleini*, *Myctophum affine*, *M. nitidulum*, and *M. punctatum* (Table 2 for  $eDW$ ; Table S1 for  $WW$  and Table S2 for  $eWW$ , Supplementary material). The 30.5% species showed an allometric coefficient higher for LWRs fitting with  $eDW$  in comparison with  $WW$ , and 27.8% in relation to  $eWW$ . Nevertheless, 52.8% and 59.3% of cases provided similar  $b$  values ( $\pm 0.1$ ). In general, the greatest differences between the growth patterns were observed for 6 myctophids: *Lampanyctus pusillus*, *L. alatus*, *Diaphus metopoclampus*, *Benthosema glaciale*, *Ceratoscopelus warmingii* and *Diaphus holti*; and for the gonostomatid *Cyclothone pseudopallida*. In these cases, the higher growth rates ( $b$ ) of the RLWs were estimated for  $eDW$ . In contrast, a negative difference between  $b$  values was noted for: *Sigmops elongatus* (Gonostomatidae), *Notolychnus valdiviae* (Myctophidae), *Argyropelecus sladeni* (Sternoptychidae), and *Chauliodus danae* (Stomiidae). Strong correlations were found between the values of  $b$  and  $\log a$ , varying significantly between fish body shapes ( $F_{2,32} = 3.486$ ,  $P = 0.044$ , Figure 2). The slope was larger in elongated shapes ( $b = -1.878$ ) and smaller in short and compressed fishes as sternoptychids ( $b = -1.056$ ).

Intra-specific differences between growth patterns within species were higher when comparing  $eWW$  and  $eDW$  than between  $WW$  and  $eWW$ , which revealed the important contribution of water content. The species with higher water content ( $> 80\%$ ) were *Cyclothone pallida*, *C. acclinidens*, *Lampanyctus nobilis*, *Chauliodus danae*, *Sigmops elongatus* and

*Stomias boa boa* (Table 3). By contrast, *Notolychnus valdiviae*, *Polyipnus polli*, *Lampanyctus pusillus*, *Benthoosema glaciale* and *Diaphus fragilis* had values lower than 75% (Table 3). The elongated shaped fishes were characterized by higher water content (mean and standard deviation,  $81.50 \pm 1.85$ ) than fusiform fishes ( $76.41 \pm 4.11$ ), and the short and deep shaped species ( $75.99 \pm 4.06$ ). A high variability was found between groups (Kruskal-Wallis test,  $\chi^2 = 12.81$ ,  $df = 3$ ,  $P = 0.002$ ).

The mesopelagic fishes showed similar average values of  $K_{rel}$  ranging between 0.792 (*Diaphus brachycephalus* from the oligotrophic region) and 1.120 (*D. rafinesquii* from the Cape Blanc upwelling region). Regional associations in the average values of  $K_{rel}$  were found for *D. brachycephalus* ( $t$ -test,  $t = 8.817$ ,  $df = 84$ ,  $P < 0.0001$ ) with lower values ( $0.792 \pm 0.185$ ) in oligotrophic waters than in equatorial region ( $0.902 \pm 0.076$ ) (Table 4), and in *Lampanyctus alatus* (Kruskal-Wallis test,  $\chi^2 = 17.309$ ,  $df = 3$ ,  $P = 0$ ), showing an increase of average  $K_{rel}$  for regions closer to African coast. The equatorial region differed from the Cape Blanc upwelling region and Canary Islands, but it was not dissimilar to Cape Verde Islands region. The  $K_{rel}$  value of Cape Verde Islands only presented differences with the average of the Canary Islands; and the Cape Blanc upwelling region and Canary Islands also reached similar average of condition (Table 4).

#### 4. DISCUSSION

The present study contributes to knowledge of mesopelagic fishes by reporting on LWR equations for 36 species, thereby establishing the effect of different weights in the allometry of LWR's, and by comparing the environmental effect in the relative condition factor. In general, our findings seem to reinforce the theory that the growth pattern is a feature identifying each

species (Mayrat, 1970) since 24 of the 36 of mesopelagic fishes analysed demonstrated similar allometric relationships, independent of the weight measure used. Some species reached a higher  $b$  value when fitting  $WW$  instead of  $eWW$ , reflecting the influence of full guts and, to a lesser extent, gonadal mass (e.g., *Vinciguerria nimbaria*, *Chauliodus danae* and *Cyclothone livida*), but in other species showed the opposite effect (e.g., *Lobianchia dofleini*, *Lepidophanes guentheri* and *Diaphus vanhoeffeni*). Nevertheless, the use of eviscerated dry weight is always recommended for LWR's, since it more accurately reflects better the muscular growth, irrespective of the trophic behaviour (full or empty guts) or the gonadal weight (important at maturation) (Pauly 1984; Froese, 2006). In other instances, such as for reconstruction and estimation of biomass of partially digested prey in studies of trophic ecology, information on  $WW$  would also be relevant. Allometric coefficients ( $b$  for  $eDW$ ) in the present study were within the expected range (2.5-3.5) for fishes (Froese, 2006), mostly ranging between 2.952 (25% percentile) and 3.384 (75% percentile). The most atypical allometric coefficient was observed for the myctophid *Lampanyctus pusillus*, 4.247, which could be due to the small size range analysed.

The species and sizes ranges of myctophids and hatchetfishes studied here showed a positive allometric growth, which implies a faster growth in body mass than in body length, i.e., more robust body and with a greater amount of muscle mass. This feature may help in the daily feeding vertical migrations, to the surface in myctophids and to the shallower mesopelagic depths in hatchetfishes (Olivar *et al.*, 2012, 2016, 2017). The opposite pattern with negative allometric growths, i.e., faster growth in length than in weight, was mostly observed in stomiiform species such as *Chauliodus danae*, *Cyclothone acclinidens*, *C. livida*, *C. pallida* and *Sigmops elongatus*, which live in deeper waters, and which have an elongate shape and higher water content. This finding is in accordance with previous studies showing an association between the proportion of water content and distribution depth of these fishes (Childres &

Nygaard, 1973; Neighbors & Nafpaktitis, 1982; Bailey & Robison, 1986; Stickney & Torres, 1989; Pakhomov *et al.*, 1996), and is also accompanied by a decrease in skeletal ash content (see Childres & Nygaard, 1973). The main reason of this somatic growth is probably due to absence of extensive vertical migrations in most of these species (Badcock and Merret, 1976; Olivar *et al.*, 2017). In general, the allometric pattern found in our species did not differ from those in other studies, although slight variations can be due to exogenous and endogenous factors (e.g., Kimmerer *et al.*, 2005; Jobling, 2008; Mazumder *et al.*, 2016), as well as the type of length measurement (standard or total), and size-range analysed. For instance, Fock & Ehrich (2010) gave a wide list of LWR's for mesopelagic fishes from the North Atlantic. Their estimation of allometric coefficients were smaller than in our study (for *eDW*) for those species differing in the size range, for example in *B. glaciale* (3.647 vs 3.020), *D. dumerilli* (3.076 vs 3.018), *D. metopoclampus* (3.684 vs 3.074), *L. dofleini* (3.448 vs 2.609) and *N. bolini* (3.212 vs 2.331). However, they were similar when the size ranges were similar as for example in *D. holti* (3.356 vs 3.350), *M. punctatum* (3.363 vs 3.448) and *S. boa boa* (3.081 vs 3.184). For fishes with similar fish size-ranges, our allometric patterns were larger than in other geographical areas, for example in the Mediterranean Sea for *Diaphus holti* (3.356 vs 3.102) and *Lampanyctus pusillus* (4.247 vs 2.296) (Battaglia *et al.*, 2010), and in the North-western Pacific Ocean for *Ceratoscopelus warmingii* (3.537 vs 3.153) (Wang *et al.*, 2018).

Food availability and physical factors have a strong influence on the growth and condition factors (Le Cren, 1951). Olivar *et al.* (2017) found higher environmental gradients in terms of temperature (22°C of difference) and salinity (3 PSU) between the sea surface and deep waters in the subtropical western Atlantic Ocean than eastern zone (10°C and 1 PSU, respectively). Although this environmental variation may affect the energetic balance and growth of specimens, any difference was detected in most species, except for *Lampanyctus alatus*. The most probable reason for the increase of the  $K_{rel}$  along the Atlantic transect may be related to

the visual system, a more generalist pattern being characterized by a major visual field favouring the detection of preys in all directions (de Buserrolles *et al.*, 2014, de Buserrolles & Marshall, 2107), and which is more numerous in the western sector. Finally, low values of  $K_{rel}$  ( $< 1$ ) for some species (e.g., *Chauliodus danae*, *Diaphus brachycephalus*, *D. mollis* and *Vingiguerria. nimbaria*) were noted in all regions, which suggests two hypotheses: a) the energetic cost of diel vertical migration (DVM) may be higher in these species; or b) the energetic requirement may be less. Unfortunately, information on the food conversion efficiency on fish growth is not available for many mesopelagic fish species. However, the intraspecific and regional variability found in our study reinforces the importance of investigations into ecological and energetic demands of deep-sea organisms (Siebel & Drazen, 2007; Irigoien *et al.*, 2014; Belcher *et al.*, 2019).

## Acknowledgments

The authors are very grateful to the captain, crew and technicians on board the R/V Hesperides for their help during the survey, and to all colleagues who participated in the survey for their cooperation. This research was sponsored by the Spanish Ministerio de Economía y Competitividad through the project CTM2012-39587-C04-03.

## Contributions

MPO and PAH were responsible for the species identifications; CLP, conceived the initial idea and wrote the main paper in collaboration with the other co-authors; CLP, MPO and VTA conducted the statistical analysis. All authors discussed results and implications, providing significant inputs to the manuscript at all stages.

308

## 309 **References**

- 310 Atkinson, D. B. (1989). Weight-length relationships of roundnose grenadier (*Coryphaenoides rupestris*  
311 Gunn.) in different areas of the North Atlantic. *Fisheries Research*, **7**, 65-72.
- 312 Badcock, J., & Merrett, N. R. (1976). Midwater fishes in the eastern North Atlantic—I. Vertical  
313 distribution and associated biology in 30N, 23W, with developmental notes on certain  
314 myctophids. *Progress in Oceanography*, **7**, 3-58.
- 315 Bailey, T. G., & Robison, B. H. (1986). Food availability as a selective factor on the chemical  
316 compositions of midwater fishes in the eastern North Pacific. *Marine Biology*, **91**, 131-141.
- 317 Battaglia, P., Malara, D., Romeo, T., & Andaloro, F. (2010). Relationships between otolith size and fish  
318 size in some mesopelagic and bathypelagic species from the Mediterranean Sea (Strait of Messina,  
319 Italy). *Scientia Marina*, **74**, 605-612.
- 320 Battaglia, P., Malara, D., Ammendolia, G., Romeo, T., & Andaloro, F. (2015). Relationships between  
321 otolith size and fish length in some mesopelagic teleosts (Myctophidae, Paralepididae,  
322 Phosichthyidae and Stomiidae). *Journal of Fish Biology*, **87**, 774-782.
- 323 Battaglia, P., Andaloro, F., Esposito, V., Granata, A., Gugliemo, L., Gugliemo, R., Musolino, S.,  
324 Romeo, T., & Zagami, G. (2016). Diet and trophic ecology of the lanternfish *Electrona risso* (Cocco  
325 1829) in the Strait of Messina (central Mediterranean Sea) and potential resource utilization from the  
326 Deep Scattering Layer (DSL). *Journal of Marine Systems*, **159**, 100-108.
- 327 Belcher, A., Saunders, R. A., & Tarling, G. A. (2019). Respiration rates and active carbon flux of  
328 mesopelagic fishes (Family Myctophidae) in the Scotia Sea, Southern Ocean. *Marine Ecology*  
329 *Progress Series*, **610**, 149-162.
- 330 Bernal, A., Olivar, M. P., Maynou, F., & de Puellas, M. L. F. (2015). Diet and feeding strategies of  
331 mesopelagic fishes in the western Mediterranean. *Progress in Oceanography*, **135**, 1-17.
- 332 Beyer, J. E. (1987). On length-weight relationships. Part 1. Corresponding the mean weight of a given  
333 length class. *Fishbytes*, **5**, 11-13.

334 Brosset, P., Fromentin, J. M., Van Beveren, E., Lloret, J., Marques, V., Basilone, G., ... & De Felice, A.  
 335 (2017). Spatio-temporal patterns and environmental controls of small pelagic fish body condition  
 336 from contrasted Mediterranean areas. *Progress in Oceanography*, **151**, 149-162.

337 Childress, J. J., & Nygaard, M. H. (1973). The chemical composition of midwater fishes as a function  
 338 of depth of occurrence off southern California. *Deep Sea Research*, **20**, 1093-1109.

339 Christensen, V., & Pauly, D. (1992). ECOPATH II—A software for balancing steady-state ecosystem  
 340 models and calculating network characteristics. *Ecological modelling*, **61**, 169-185.

341 Christensen, V., & Walters, C. J. (2004). Ecopath with Ecosim: methods, capabilities and  
 342 limitations. *Ecological modelling*, **172**, 109-139.

343 Choy, C. A., Popp, B. N., Hannides, C. C. S., & Drazen, J. C. (2015). Trophic structure and food  
 344 resources of epipelagic and mesopelagic fishes in the North Pacific Subtropical Gyre ecosystem  
 345 inferred from nitrogen isotopic compositions. *Limnology and Oceanography*, **60**, 1156–1171.

346 Davison, P. C., Checkley Jr., D. M., Koslow, J. A., & Barlow, J. (2013). Carbon export mediated by  
 347 mesopelagic fishes in the northeast Pacific Ocean. *Progress in Oceanography*, **116**, 14-30.

348 De Busserolles, F., Marshall, N. J., & Collin, S. P. (2014). The eyes of lanternfishes (Myctophidae,  
 349 teleostei): novel ocular specializations for vision in dim light. *Journal of Comparative*  
 350 *Neurology*, **522**, 1618-1640.

351 De Busserolles, F., & Marshall, N. J. (2017). Seeing in the deep-sea: visual adaptations in  
 352 lanternfishes. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **372**,  
 353 20160070.

354 Dinno, A. (2017). dunn.test: Dunn's test of multiple comparisons using rank sums. R package version  
 355 1.4.

356 Drazen, J. C., & Sutton, T. T., (2017). Dining in the deep: the feeding ecology of deep-sea fishes. *Annual*  
 357 *Review of Marine Science* **9**, 337–366.

358 Eduardo, L. N., Mincarone, M. M., Villarins, B. T., Frédou, T., Lira, A. S., Bertrand, A., & Lucena-  
 359 Frédou, F. (2019). Length–weight relationships of eleven mesopelagic fishes from oceanic islands  
 360 of the Southwestern Tropical Atlantic. *Journal of Applied Ichthyology*, **35**, 605-607.

361 Efitre, J., Chapman, L. J., & Murie, D. J. (2009). Fish condition in introduced tilapias of Ugandan crater  
362 lakes in relation to deforestation and fishing pressure. *Environmental Biology of Fishes*, **85**, 63-75.

363 Ekau, W., Auel, H., Pörtner, H. O., & Gilbert, D. (2010). Impacts of hypoxia on the structure and  
364 processes in pelagic communities (zooplankton, macro-invertebrates and fish). *Biogeosciences*, **7**,  
365 1669-1699.

366 Froese, R. (2006). Cube law, condition factor and weight–length relationships: history, meta-analysis  
367 and recommendations. *Journal of Applied Ichthyology*, **22**, 241-253.

368 Fock, H. O., & Ehrich, S. (2010). Deep-sea pelagic nekton biomass estimates in the North Atlantic:  
369 horizontal and vertical resolution of revised data from 1982 and 1983. *Journal of Applied*  
370 *Ichthyology*, **26**, 85-101.

371 Fock, H. O., & Czudaj, S. (2018). Size structure changes of mesopelagic fishes and community biomass  
372 size spectra along a transect from the equator to the Bay of Biscay collected in 1966–1979 and 2014–  
373 2015. *ICES Journal of Marine Science*, **76**, 755-770.

374 Fulton, T. W. (1911). *The sovereignty of the sea: an historical account of the claims of England to the*  
375 *dominion of the British seas and of the evolution of the territorial waters, with special reference to*  
376 *the rights of fishing and the naval salute*. Edinburgh: W. Blackwood.

377 Gjøsaeter, J., & Kawaguchi, K. (1980). *A review of the world resources of mesopelagic fish*. Fish FAO  
378 *Fisheries Technical Paper*, **193**, 1-151.

379 Gould, S. J. (1966). Allometry and size in ontogeny and phylogeny. *Biological Reviews*, **41**, 587-638.

380 Hudson, J. M., Steinberg, D. K., Sutton, T. T., Graves, J. E., & Latour, R. J. (2014). Myctophid feeding  
381 ecology and carbon transport along the northern Mid-Atlantic Ridge. *Deep Sea Research Part I:*  
382 *Oceanographic Research Papers*, **93**, 104-116.

383 Hulley, P. A. (1981). Results of the research cruises of FRV" Walther Herwig" to South America LVIII,  
384 Family Myctophidae (Osteichthyes, Myctophiformes). *Archiv für Fischereiwissenschaft*, **31**, 1-  
385 300.

386 Irigoien, X., Klevjer, T. A., Røstad, A., Martinez, U., Boyra, G., Acuña, J. L., ... & Agusti, S. (2014).  
387 Large mesopelagic fishes biomass and trophic efficiency in the open ocean. *Nature*  
388 *communications*, **5**, 3271.



Ikeda, T., Shiga, N., & Yamaguchi, A. (2008). Structure, biomass distribution and trophodynamics of the pelagic ecosystem in the Oyashio region, western subarctic Pacific. *Journal of Oceanography*, **64**, 339-354.

Jellyman, P. G., Booker, D. J., Crow, S. K., Bonnett, M. L., & Jellyman, D. J. (2013). Does one size fit all? An evaluation of length–weight relationships for New Zealand's freshwater fish species. *New Zealand Journal of Marine and Freshwater Research*, **47**, 450-468.

Jiang, Y. E., Chen, Z. Z., Zhang, K., Zhang, J., Gong, Y. Y., Kong, X. L., ... & Fang, Z. Q. (2017). Length–weight relationships of seven myctophid fishes (Myctophiformes: Myctophidae) in the South China Sea. *Journal of Applied Ichthyology*, **33**, 1044-1046.

Jobling, M. (2008). Environmental factors and rates of development and growth. In P. J. Hart & J. D. Reynolds (Eds.), *Handbook of fish biology and fisheries, Vol. 1: fish biology* (pp. 97–122). Oxford: Blackwell Publishing Ltd.

Kimmerer, W., Avent, S. R., Bollens, S. M., Feyrer, F., Grimaldo, L. F., Moyle, P. B., Nobriga, M., & Visintainer, T. (2005) Variability in length–weight relationships used to estimate biomass of estuarine fish from survey data. *Transactions of the American Fisheries Society*, **134**, 481-495.

Kulbicki, M., Guillemot, N., & Amand, M. (2005). A general approach to length-weight relationships for New Caledonian lagoon fishes. *Cybium*, **29**, 235-252.

Krefft, G. (1976). Distribution patterns of oceanic fishes in the Atlantic Ocean. Selected problems. *Revue des Travaux de l'Institut des Pêches Maritimes*, **40**, 439-460.

Le Cren, E. D. (1951). The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *The Journal of Animal Ecology*, **20**, 201-219.

Lloret, J., Gil de Sola, L., Souplet, A., & Galzin, R. (2002). Effects of large-scale habitat variability on condition of demersal exploited fish in the north-western Mediterranean. *ICES Journal of Marine Science*, **59**, 1215-1227.

Lloret, J., Shul'man, G. E., & Love, R. M. (2014). *Condition and health indicators of exploited marine fishes*. Oxford: Wiley Blackwell.

Margalef, R. (1974). *Ecologia*. Barcelona: Ediciones Omega.

Mayrat, A. (1970). Allometrie et taxinomie. *Revue de Statistique Appliquée*, **18**, 47-58.

417 Mazumder, S. K., Das, S. K., Bakar, Y., & Ghaffar, M. A. (2016). Effects of temperature and diet on  
 418 length-weight relationship and condition factor of the juvenile Malabar blood snapper (*Lutjanus*  
 419 *malabaricus* Bloch & Schneider, 1801). *Journal of Zhejiang University-SCIENCE B*, **17**, 580-590.  
 420 Millar, J. S., & Hickling, G. J. (1990). Fasting endurance and the evolution of mammalian body  
 421 size. *Functional Ecology*, **4**, 5-12.  
 422 Mittelstaedt, E. (1983). The upwelling area off Northwest Africa—a description of phenomena related  
 423 to coastal upwelling. *Progress in Oceanography*, **12**, 307-331.  
 424 Moffitt, S. E., Hill, T. M., Ohkushi, K., Kennett, J. P., & Behl, R. J. (2014). Vertical oxygen minimum  
 425 zone oscillations since 20 ka in Santa Barbara Basin: A benthic foraminiferal community  
 426 perspective. *Paleoceanography*, **29**, 44-57.  
 427 Morel, A., Claustre, H., & Gentili, B. (2010). The most oligotrophic subtropical zones of the global  
 428 ocean: similarities and differences in terms of chlorophyll and yellow substance. *Biogeosciences*, **7**,  
 429 3139-3151.  
 430 Neighbors, M. A., & Nafpaktitis, B. G. (1982). Lipid compositions, water contents, swimbladder  
 431 morphologies and buoyancies of nineteen species of midwater fishes (18 myctophids and 1  
 432 neoscopelid). *Marine Biology*, **66**, 207-215.  
 433 Nelson, J. S., Grande, T. C., & Wilson, M. V. H. (2016). *Fishes of the World*. Fifth Edition. New Jersey:  
 434 John Wiley & Sons, Inc.  
 435 Olivar, M.P., Bernal, A., Molí, B., Peña, M., Balbín, R., Castellón, A., Miquel, J., & Massutí, E. (2012).  
 436 Vertical distribution, diversity and assemblages of mesopelagic fishes in the western Mediterranean.  
 437 *Deep-Sea Research Part I*, **62**, 53–69.  
 438 Olivar, M.P., González-Gordillo, J.I., Salat, J., Chust, G., Cózar, A., Hernández-León, S., Fernández de  
 439 Puellas, M.L., & Irigoien, X. (2016). The contribution of migratory mesopelagic fishes to neuston  
 440 fish assemblages across the Atlantic, Indian and Pacific Oceans. *Marine and Freshwater Research*,  
 441 **67**, 1114–1127.  
 442 Olivar, M. P., Hulley, P. A., Castellón, A., Emelianov, M., López, C., Tuset, V. M., ... & Molí, B. (2017).  
 443 Mesopelagic fishes across the tropical and equatorial Atlantic: biogeographical and vertical  
 444 patterns. *Progress in oceanography*, **151**, 116-137.

445   Olivar, M. P., Bode, A., López-Pérez, C., Hulley, P. A., & Hernández-León, S. (2018). Trophic position  
446   of lanternfishes (Pisces: Myctophidae) of the tropical and equatorial Atlantic estimated using stable  
447   isotopes. *ICES Journal of Marine Science*, **76**, 649-661.

448   Pakhomov, E. A., Perissinotto, R., & McQuaid, C. D. (1996). Prey composition and daily rations of  
449   myctophid fishes in the Southern Ocean. *Marine Ecology Progress Series*, **134**, 1-14.

450   Pauly, D. (1984). Fish population dynamics in tropical waters: a manual for use with programmable  
451   calculators. *ICLARM Studies and Reviews*, **8**, 1-325.

452   Pauly, D. (1993). Special Book Section: Data-Rich Books. *BioScience*, **43**, 167-168.

453   Pauly, D. (2000). Global change, fisheries, and the Integrity of Marine Ecosystems; the future has  
454   already begun. In D. Pimentel, L. Westra & R.F. Ross (Eds.), *Ecological Integrity: Integrating*  
455   *Environment, Conservation, and Health* (pp. 227-239). Washington; Islands Press.

456   Pazianoto, L. H., Cionek, V. D. M., Fogaça, F. N., & Benedito, E. (2016). Feeding and body condition  
457   of an invasive fish species under different environmental conditions. *Neotropical Ichthyology*, **14**,  
458   e150164.

459   Pereira, J. N., Neves, V. C., Prieto, R., Silva, M. A., Cascão, I., Oliveira, C., ... & Clarke, D. (2011).  
460   Diet of mid-Atlantic Sowerby's beaked whales *Mesoplodon bidens*. *Deep Sea Research Part I:*  
461   *Oceanographic Research Papers*, **58**, 1084-1090.

462   R Core Team, 2016. R: a Language and Environment for Statistical Computing. R Foundation for  
463   Statistical Computing, Vienna, Austria. <http://www.r-project.org>.

464   Sakuma, K. M., Ralston, S., Lenarz, W. H., Embury, M. (1999). Effects of the parasitic copepod  
465   *Cardiodectes medusaeus* on the lanternfishes *Diaphus theta* and *Tarletonbeania crenularis* off  
466   Central California. *Environmental Biology of Fishes* **55**, 423–430.

467   Schneider, J. C., Laarman, P. W., Howard Gowing, J. C., Laarman, P. W., & Gowing, H. (2000). Length-  
468   Weight Relationships. In J. C. Schneider (Ed.), *Manual of Fisheries Survey Methods II* (pp. 17.1–  
469   17.16), Michigan: Michigan Department of Natural Resources.

470   Seibel, B. A. & Drazen, J. C., (2007). Depth-related trends in metabolism of benthic and benthopelagic  
471   deep-sea fishes. *Limnology and Oceanography*, **52**, 2306-2316.

472 Simpkins, D. G., & Hubert, W. A. (2000). Drifting invertebrates, stomach contents, and body conditions  
 473 of juvenile rainbow trout from fall through winter in a Wyoming tailwater. *Transactions of the*  
 474 *American Fisheries Society*, **129**, 1187-1195.

475 Smith, A. D., Brown, C. J., Bulman, C. M., Fulton, E. A., Johnson, P., Kaplan, I. C., ... & Shin, Y. J.  
 476 (2011a). Impacts of fishing low-trophic level species on marine ecosystems. *Science*, **333**, 1147-  
 477 1150.

478 Sokal, R. R., Rohlf, F. J., & Lahoz León, M. (1979). *Biometry: the principles and practice of statistics*  
 479 *in biological research*. San Francisco: W.H. Freeman.

480 Springer, A. M., Piatt, J. F., Shuntov, V. P., Van Vliet, G. B., Vladimirov, V. L., Kuzin, A. E., & Perlov,  
 481 A. S. (1999). Marine birds and mammals of the Pacific Subarctic Gyres. *Progress in*  
 482 *Oceanography*, **43**, 443-487.

483 Stevenson, R. D., & Woods Jr, W. A. (2006). Condition indices for conservation: new uses for evolving  
 484 tools. *Integrative and Comparative Biology*, **46**, 1169-1190.

485 Stickney, D. G., & Torres, J. J. (1989). Proximate composition and energy content of mesopelagic fishes  
 486 from the eastern Gulf of Mexico. *Marine Biology*, **103**, 13-24.

487 Stramma, L., & Schott, F. (1999). The mean flow field of the tropical Atlantic Ocean. *Deep Sea*  
 488 *Research Part II: Topical Studies in Oceanography*, **46**, 279-303.

489 Sutton, T. T., Porteiro, F. M., Heino, M., Byrkjedal, I., Langhelle, G., Anderson, C. I. H., ... & Bergstad,  
 490 O. A. (2008). Vertical structure, biomass and topographic association of deep-pelagic fishes in  
 491 relation to a mid-ocean ridge system. *Deep Sea Research Part II: Topical Studies in*  
 492 *Oceanography*, **55**, 161-184.

493 Sutton, T. T. (2013). Vertical ecology of the pelagic ocean: classical patterns and new  
 494 perspectives. *Journal of Fish Biology*, **83**, 1508-1527.

495 Torres, M. A., Ramos, F., & Sobrino, I. (2012). Length-weight relationships of 76 fish species from the  
 496 Gulf of Cadiz (SW Spain). *Fisheries Research*, **127**, 171-175.



497 Trueman, C. N., Johnston, G., O'Hea, B., & MacKenzie, K. M. (2014). Trophic interactions of fish  
 498 communities at midwater depths enhance long-term carbon storage and benthic production on  
 499 continental slopes. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20140669.

- Verdiell-Cubedo, D., Oliva-Paterna, F. J., & Torralva, M. (2006). Length–weight relationships for 22 fish species of the Mar Menor coastal lagoon (western Mediterranean Sea). *Journal of Applied Ichthyology*, **22**, 293-294
- Wang, L., Li, Y., Zhang, R., Tian, Y., & Lin, L. (2018). Length-weight relationships of five lanternfishes (Myctophidae) from the high seas of Northwestern Pacific Ocean. *Journal of Applied Ichthyology*, **34**, 1340-1342.
- Watanabe, H., & Kawaguchi, K. (2003). Decadal change in the diets of the surface migratory myctophid fish *Myctophum nitidulum* in the Kuroshio region of the western North Pacific: Predation on sardine larvae by myctophids. *Fisheries Science*, **69**, 716-721.
- Wege, G. J., & Anderson, R. O. (1978). Relative weight (Wr): a new index of condition for largemouth bass. New approaches to the management of small impoundments. *American Fisheries Society, North Central Division, Special Publication*, **5**, 79-91.
- Wilson, A. J., & Nussey, D. H. (2010). What is individual quality? An evolutionary perspective. *Trends in Ecology & Evolution*, **25**, 207-214.

## Legends

**FIGURE 1** Location of the stations sampled during the survey across the equatorial and tropical Atlantic during April 2015, and oceanographic regions according to [Olivar et al. \(2017\)](#).

**FIGURE 2** Relationships between the regression coefficients  $\log a$  and  $b$  of LWR equation for different body shapes (short-deep, elongate and fusiform) at the species level.



**TABLE 1** List of mesoplegic fishes from equatorial and tropical Atlantic waters analyzed in the present study. Region indicates the sampling origin according to Olivar *et al.* (2017): CB, Cape Blanc; CI, Canary Islands; CV, Cape Verde; O, oligotrophic; E, equatorial. Taxa were ordered according to Nelson *et al.* (2016)

Order	Family	Genera	Species	Author	Region	
Stomiiformes	Gonostomatidae	<i>Cyclothone</i>	<i>Cyclothone acclinidens</i>	Garman, 1899	CV, CB	
			<i>Cyclothone livida</i>	Brauer, 1902	CV, CB	
			<i>Cyclothone pallida</i>	Brauer, 1902	CV, CB	
			<i>Cyclothone pseudopallida</i>	Mukhacheva, 1964	CV	
	Phosichthyidae	<i>Gonostoma</i>	<i>Sigmops elongatus</i>	(Günther, 1878)	E, CV, CB, CI	
		<i>Vinciguerrria</i>	<i>Vinciguerrria nimbaria</i>	(Jordan & Williams, 1895)	CV, CB	
	Sternoptychidae	<i>Argyropelecus</i>	<i>Argyropelecus sladeni</i>	Regan, 1908	E , CV, CB	
		<i>Polypnus</i>	<i>Polyipnus polli</i>	Schultz, 1961	CV, CB	
		<i>Sternoptyx</i>	<i>Sternoptyx diaphana</i>	Hermann, 1781	CV, CB	
	Stomiidae	<i>Chauliodus</i>	<i>Chauliodus danae</i>	Regan & Trewavas, 1929	E, CV, CB	
		<i>Stomias</i>	<i>Stomias boa boa</i>	(Risso, 1810)	CV	
Myctophiformes	Myctophidae	<i>Benthoosema</i>	<i>Benthoosema glaciale</i>	(Reinhardt, 1837)	CB	
			<i>Benthoosema suborbitale</i>	(Gilbert, 1913)	CV, CI	
		<i>Bolinichthys</i>	<i>Bolinichthys photothorax</i>	(Parr, 1928)	O, E, CV	
		<i>Ceratoscopelus</i>	<i>Ceratoscopelus warmingii</i>	(Lütken, 1892)	O, E, CV	
		<i>Diaphus</i>	<i>Diaphus brachycephalus</i>	Tåning, 1928	O, E , CV, CB	
			<i>Diaphus dumerilii</i>	(Bleeker, 1856)	E	
			<i>Diaphus fragilis</i>	Tåning, 1928	E	
			<i>Diaphus holti</i>	Tåning, 1918	CV, CB	
			<i>Diaphus metopoclampus</i>	(Cocco, 1829)	E, CI	
			<i>Diaphus mollis</i>	Tåning, 1928	O, E, CV	
			<i>Diaphus problematicus</i>	Parr, 1928	E	
			<i>Diaphus rafinesquii</i>	(Cocco, 1838)	CB, CI	
			<i>Diaphus vanhoeffeni</i>	(Brauer, 1906)	CV	
			<i>Hygophum</i>	<i>Hygophum macrochir</i>	(Günther, 1864)	CB (30)
		<i>Lampanyctus</i>	<i>Lampanyctus alatus</i>	Goode & Bean, 1896	E , CV, CB, CI	
			<i>Lampanyctus nobilis</i>	Tåning, 1928	O, E	
			<i>Lampanyctus pusillus</i>	(Johnson, 1890)	E, CB, CI	
		<i>Lepidophanes</i>	<i>Lepidophanes guentheri</i>	(Goode & Bean, 1896)	O, E, CV	
		<i>Lobianchia</i>	<i>Lobianchia dofleini</i>	(Zugmayer, 1911)	CV, CB, CI	
		<i>Myctophum</i>	<i>Myctophum affine</i>	(Lütken, 1892)	CV	
			<i>Myctophum nitidulum</i>	Garman, 1899	CV, CI	
			<i>Myctophum punctatum</i>	Rafinesque, 1810	CB	
			<i>Notolychnus</i>	<i>Notolychnus valdiviae</i>	(Brauer, 1904)	E, CV
			<i>Notoscopelus</i>	<i>Notoscopelus bolini</i>	Nafpaktitis, 1975	CB
			<i>Notoscopelus resplendens</i>	(Richardson, 1845)	E, CV, CB, CI	

528

529

530

531

532

**TABLE 2** Length-weight parametres for mesopelagic species (ordered alphabetically) from the equatorial and tropical Atlantic waters. *a*, intercept; *a*+, positive allometry; *a*-, negative allometry; *b*, allometry coefficient (slope); *eDW*, eviscerate body dry weight; *i*, isometry; *n*, sample size; *SL*, standard length; *r*<sup>2</sup>, coefficient of determination; 95% CL of *b*, confidence interval

Species	n	Range <i>SL</i> (mm)	Range <i>eDW</i> (g)	<i>a</i>	<i>b</i>	Inferior 95% CL of <i>b</i>	Superior 95% CL of <i>b</i>	<i>r</i> <sup>2</sup>	Growth model
<i>Argyropelecus sladeni</i>	27	15 – 43	0.012 – 0.186	0.00000414	2.950	2.472	3.428	0.866	i
<i>Benthoosema glaciale</i>	15	15 – 35	0.005 – 0.145	0.00000035	3.647	3.486	3.809	0.990	a+
<i>Benthoosema suborbitale</i>	29	16 – 33	0.012 – 0.103	0.00000198	3.135	2.868	3.401	0.955	i
<i>Bolinichthys photothorax</i>	6	21 – 51	0.021 – 0.335	0.00000121	3.183	2.944	3.421	0.997	i
<i>Ceratoscopelus warmingii</i>	47	16 – 64	0.006 – 0.796	0.00000032	3.537	3.443	3.631	0.992	a+
<i>Chauliodus danae</i>	43	25 – 212	0.023 – 3.519	0.00000139	2.710	2.554	2.867	0.968	a-
<i>Cyclothone acclinidens</i>	22	31 – 46	0.005 – 0.051	0.00000524	2.337	1.849	2.825	0.768	a-
<i>Cyclothone livida</i>	32	23 – 37	0.007 – 0.030	0.00000372	2.477	2.011	2.941	0.798	a-
<i>Cyclothone pallida</i>	24	25 – 54	0.009 – 0.073	0.00000235	2.600	2.270	2.929	0.924	a-
<i>Cyclothone pseudopallida</i>	19	23 – 35	0.004 – 0.017	0.00000258	2.420	1.596	3.244	0.693	i
<i>Diaphus brachycephalus</i>	100	12 – 47	0.002 – 0.749	0.00000131	3.339	3.194	3.482	0.956	a+
<i>Diaphus dumerilii</i>	30	31 – 57	0.089 – 0.628	0.00000254	3.076	2.865	3.285	0.970	i
<i>Diaphus fragilis</i>	10	46 – 79	0.183 – 1.022	0.00000155	3.056	2.740	3.371	0.984	i
<i>Diaphus holti</i>	91	11 – 50	0.002 – 0.491	0.00000106	3.356	3.250	3.460	0.978	a+
<i>Diaphus metopoclampus</i>	10	19 – 40	0.009 – 0.182	0.00000022	3.684	3.219	4.147	0.977	a+
<i>Diaphus mollis</i>	30	29 – 55	0.073 – 0.571	0.00000247	3.093	2.912	3.272	0.979	i
<i>Diaphus problematicus</i>	9	45 – 72	0.223 – 0.787	0.00000437	2.820	2.441	3.199	0.978	i
<i>Diaphus rafinesquii</i>	125	11 – 70	0.004 – 1.425	0.00000242	3.107	3.042	3.170	0.987	a+
<i>Diaphus vanhoeffeni</i>	17	25 – 36	0.054 – 0.154	0.00001968	2.488	1.856	3.119	0.825	i
<i>Hygophum macrochir</i>	30	20 – 42	0.018 – 0.223	0.00000190	3.138	2.973	3.302	0.982	i
<i>Lampanyctus alatus</i>	142	16 – 57	0.006 – 0.362	0.00000023	3.594	3.492	3.696	0.972	a+
<i>Lampanyctus nobilis</i>	19	28 – 74	0.022 – 0.619	0.00000061	3.205	2.906	3.503	0.968	i
<i>Lampanyctus pusillus</i>	22	24 – 36	0.019 – 0.132	0.00000003	4.247	3.638	4.855	0.914	a+
<i>Lepidophanes guentheri</i>	35	19 – 61	0.008 – 0.521	0.00000043	3.391	3.181	3.601	0.970	a+
<i>Lobianchia dofleini</i>	30	13 – 30	0.006 – 0.118	0.00000090	3.448	3.081	3.813	0.930	a+
<i>Myctophum affine</i>	34	14 – 47	0.005 – 0.328	0.00000049	3.509	3.384	3.633	0.990	a+
<i>Myctophum nitidulum</i>	25	15 – 74	0.006 – 1.233	0.00000084	3.308	3.219	3.396	0.996	a+
<i>Myctophum punctatum</i>	28	16 – 69	0.007 – 1.068	0.00000057	3.363	3.252	3.472	0.993	a+
<i>Notolychnus valdiviae</i>	27	15 – 22	0.019 – 0.123	0.00000289	2.956	2.956	3.577	0.816	i
<i>Notoscopelus bolini</i>	11	21 – 28	0.015 – 0.040	0.00000092	3.212	2.664	3.760	0.951	i
<i>Notoscopelus resplendens</i>	20	20 – 70	0.016 – 1.085	0.00000179	3.111	2.941	3.280	0.988	i
<i>Polyipnus polli</i>	34	16 – 43	0.019 – 0.627	0.00000064	3.697	3.431	3.962	0.962	a+
<i>Sternoptyx diaphana</i>	29	9 – 41	0.002 – 0.334	0.00000277	3.054	2.879	3.228	0.978	i
<i>Stomias boa boa</i>	20	53 – 153	0.041 – 1.073	0.00000019	3.081	2.855	3.305	0.979	i
<i>Sigmops elongatus</i>	35	45 – 151	0.029 – 0.962	0.00000089	2.746	2.608	2.883	0.980	a-
<i>Vinciguerrria nimbaria</i>	32	14 – 50	0.005 – 0.226	0.00000177	3.005	2.932	3.077	0.996	i



**TABLE 3** Mean water content (and standard deviation, sd) expressed as % of eviscerated wet weight in mesopelagic fishes from the equatorial and tropical Atlantic waters

<b>Species</b>	<b>mean <math>\pm</math> sd</b>
<i>Notolychnus valdiviae</i>	58.99 $\pm$ 19.97
<i>Polyipnus polli</i>	71.31 $\pm$ 2.95
<i>Lampanyctus pusillus</i>	73.50 $\pm$ 3.73
<i>Benthoosema glaciale</i>	73.63 $\pm$ 3.59
<i>Diaphus fragilis</i>	74.67 $\pm$ 0.80
<i>Diaphus vanhoeffeni</i>	75.14 $\pm$ 1.29
<i>Myctophum nitidulum</i>	75.19 $\pm$ 2.04
<i>Diaphus problematicus</i>	75.34 $\pm$ 1.20
<i>Lobianchia dofleini</i>	75.70 $\pm$ 2.28
<i>Myctophum affine</i>	75.76 $\pm$ 2.00
<i>Diaphus dumerilii</i>	76.20 $\pm$ 1.67
<i>Vinciguerrria nimbaria</i>	76.22 $\pm$ 1.47
<i>Notoscopelus resplendens</i>	76.49 $\pm$ 1.44
<i>Lepidophanes guentheri</i>	76.50 $\pm$ 2.79
<i>Diaphus holti</i>	76.58 $\pm$ 3.44
<i>Diaphus brachycephalus</i>	76.89 $\pm$ 1.71
<i>Diaphus mollis</i>	77.49 $\pm$ 1.34
<i>Benthoosema suborbitale</i>	77.60 $\pm$ 1.47
<i>Diaphus rafinesquii</i>	77.69 $\pm$ 2.01
<i>Argyropelecus sladeni</i>	78.15 $\pm$ 2.01
<i>Lampanyctus alatus</i>	78.24 $\pm$ 2.97
<i>Sternoptyx diaphana</i>	78.51 $\pm$ 2.90
<i>Diaphus metopoclampus</i>	79.05 $\pm$ 2.86
<i>Cyclothone pseudopallida</i>	79.07 $\pm$ 2.93
<i>Bolinichthys photothorax</i>	79.37 $\pm$ 1.30
<i>Myctophum punctatum</i>	79.53 $\pm$ 2.18
<i>Cyclothone livida</i>	79.69 $\pm$ 2.02
<i>Notoscopelus bolini</i>	79.77 $\pm$ 1.13
<i>Ceratoscopelus warmingii</i>	79.84 $\pm$ 2.49
<i>Hygophum macrochir</i>	79.93 $\pm$ 1.51
<i>Cyclothone pallida</i>	80.66 $\pm$ 2.19
<i>Cyclothone acclinidens</i>	81.21 $\pm$ 2.37
<i>Lampanyctus nobilis</i>	81.54 $\pm$ 1.51
<i>Chauliodus danae</i>	82.65 $\pm$ 5.82
<i>Sigmops elongatus</i>	83.35 $\pm$ 2.42
<i>Stomias boa boa</i>	83.90 $\pm$ 1.64

**TABLE 4** Statistical comparison of Relative Condition Factor ( $K_{rel}$ ) between oceanographic regions of the study area (Olivar *et al.*, 2017) for mesopelagic fish species from the equatorial and tropical Atlantic waters. CB, Cape Blanc; CI, Canary Islands; CV, Cape Verde; E, equatorial;  $eWD$ , eviscerated dry weight (mg) ;  $LWR$ , length-weight relationship; n, number of specimens; ns, non-significant;  $KW$ , Kruskal-Wallis test; O, oligotrophic; sd, standard deviation;  $SL$ , standard length (mm); U, Mann-Whitney U test

Species	Region	n	SL range	eDW range	LWR equation	r <sup>2</sup>	K <sub>rel</sub> mean (sd)	Statistical analysis			
								test	df	P	
<i>Chauliodus dane</i>	E	12	40-176	0.024-1.758	log eWD = 2.867 log SL -6.182	0.993	0.936 (0.090)	t-test = -0.294	33	ns	
	CV	23	49-176	0.047-3.520			0.948 (0.112)				
<i>Ceratoscopelus warmingii</i>	E	23	16-64	0.071-0.796	log eWD = 3.610 log SL -6.581	0.992	1.072 (0.185)	U = 129, z = -1.552		ns	
	CV	16	36-64	0.105-0.768			0.999 (0.101)				
<i>Diaphus brachycephalus</i>	O	34	30-53	0.101-0.606	log eWD = 3.264 log SL -5.767	0.972	0.792 (0.067)	t-test = -6.817	84	< 0.0001	
	E	53	16-53	0.015-0.750			0.902 (0.076)				
<i>Diaphus mollis</i>	E	12	34-52	0.134-0.572	log eWD = 3.182 log SL -5.748	0.985	0.914 (0.076)	t-test = 1.169	20	ns	
	CV	10	29-54	0.074-0.566			0.877 (0.069)				
<i>Diaphus rafinesquii</i>	CB	90	30-70	0.101-1.425	log eWD = 3.079 log SL -5.573	0.984	1.120 (0.094)	t-test = 0.063*	47	ns	
	CI	34	30-58	0.118-0.773			1.118 (0.125)				
<i>Lepidophanes guentheri</i>	O	16	19-44	0.008-0.137	log eWD = 3.479 log SL -6.513	0.984	0.998 (0.187)	t-test = -1.312	28	ns	
	E	14	34-61	0.077-0.522			1.088 (0.186)				
<i>Myctophum nitidulum</i>	CV	10	15-74	0.101-0.238	log eWD = 3.308 log SL -6.074	0.984	1.070 (0.102)	U = 72, z =-0.166		ns	
	CI	15	18-57	0.118-0.231			1.051 (0.115)				
<i>Vinciguerrria nimbaria</i>	CV	17	37-50	0.101-0.238	log eWD = 3.005 log SL -5.752	0.996	0.891 (0.083)	t-test = 0.088	30	ns	
	CB	15	14-20	0.118-0.231			0.888 (0.083)				
								Post-hoc test after KW			
								CV	CB	CI	
<i>Lampanyctus alatus</i>	E	22	27-48	0.027-0.217	log eWD = 3.594 log SL -6.648	0.972	1.067 (0.151)	ns	0.024	0.002	
	CV	69	16-57	0.006-0.358			1.103 (0.152)				ns
	CB	25	34-51	0.072-0.363			1.195 (0.149)				
	CI	26	39-53	0.150-0.347			1.234 (0.171)				

\* $t$ -test for unequal variance

**TABLE S1** Length-weight parameters for mesopelagic species (ordered alphabetically) from the equatorial and tropical Atlantic waters. *a*, intercept; *a*+, positive allometry; *a*-, negative allometry; *b*, allometry coefficient (slope); *WW*, whole body wet weight; *i*, isometry; *n*, sample size; *SL*, standard length; *r*<sup>2</sup>, correlation of determination; 95% CL of *b*, confidence interval

Species	n	Range <i>SL</i> (mm)	Range <i>WW</i> (g)	<i>a</i>	<i>b</i>	Inferior 95% CL of <i>b</i>	Superior 95% CL of <i>b</i>	<i>r</i> <sup>2</sup>	Growth model
<i>Argyroleucus sladeni</i>	27	15 – 43	0.061 – 1.054	0.00000925	3.223	2.692	3.753	0.862	i
<i>Benthoosema glaciale</i>	15	15 – 35	0.034 – 0.597	0.00000539	3.251	3.100	3.400	0.989	a+
<i>Benthoosema suborbitale</i>	29	16 – 33	0.069 – 0.527	0.00001615	2.982	2.592	3.371	0.943	i
<i>Bolinichthys photothorax</i>	6	21 – 51	0.126 – 1.625	0.00001123	3.022	2.607	3.435	0.990	i
<i>Ceratoscopelus warmingii</i>	47	16 – 64	0.038 – 3.865	0.00000442	3.284	3.195	3.371	0.992	a+
<i>Chauliodus danae</i>	43	25 – 212	0.062 – 29.083	0.00000267	2.976	2.868	3.083	0.987	i
<i>Cyclothone acclinidens</i>	22	31 – 46	0.039 – 0.340	0.00004436	2.243	1.779	2.705	0.772	a-
<i>Cyclothone livida</i>	32	23 – 37	0.040 – 0.195	0.00000599	2.846	2.340	3.351	0.815	i
<i>Cyclothone pallida</i>	24	25 – 54	0.051 – 0.434	0.00001276	2.616	2.210	3.020	0.891	i
<i>Cyclothone pseudopallida</i>	19	23 – 35	0.030 – 0.087	0.00005320	2.016	1.356	2.675	0.710	a-
<i>Diaphus brachycephalus</i>	100	12 – 47	0.014 – 3.557	0.00000906	3.243	3.123	3.362	0.967	a+
<i>Diaphus dumerilii</i>	30	31 – 57	0.423 – 2.681	0.00001689	2.981	2.763	3.199	0.966	i
<i>Diaphus fragilis</i>	10	46 – 79	0.871 – 4.415	0.00000971	2.976	2.798	3.152	0.995	i
<i>Diaphus holti</i>	91	11 – 50	0.020 – 2.367	0.00001677	3.006	2.887	3.124	0.966	i
<i>Diaphus metopoclampus</i>	10	19 – 40	0.066 – 0.943	0.00000398	3.353	2.914	3.791	0.975	i
<i>Diaphus mollis</i>	30	29 – 55	0.378 – 2.883	0.00001912	2.972	2.814	3.129	0.982	i
<i>Diaphus problematicus</i>	9	45 – 72	0.971 – 3.568	0.00001890	2.834	2.498	3.168	0.983	i
<i>Diaphus rafinesquii</i>	125	11 – 70	0.027 – 6.080	0.00003096	2.850	2.788	2.913	0.985	a-
<i>Diaphus vanhoeffeni</i>	17	25 – 36	0.265 – 0.691	0.00007869	2.530	2.088	2.972	0.908	a-
<i>Hygophum macrochir</i>	30	20 – 42	0.115 – 1.192	0.00002150	2.926	2.772	3.079	0.982	i
<i>Lampanyctus alatus</i>	142	16 – 57	0.047 – 1.795	0.00000488	3.199	3.106	3.290	0.971	a+
<i>Lampanyctus nobilis</i>	19	28 – 74	0.150 – 3.349	0.00000431	3.167	2.945	3.388	0.982	i
<i>Lampanyctus pusillus</i>	22	24 – 36	0.085 – 0.517	0.00000136	3.573	3.063	4.081	0.915	a+
<i>Lepidophanes guentheri</i>	35	19 – 61	0.056 – 2.564	0.00000642	3.108	2.932	3.282	0.975	i
<i>Lobianchia dofleini</i>	30	13 – 30	0.036 – 0.508	0.00001128	3.130	2.811	3.449	0.935	i
<i>Myctophum affine</i>	34	14 – 47	0.025 – 1.910	0.00000371	3.375	3.109	3.640	0.955	a+
<i>Myctophum nitidulum</i>	25	15 – 74	0.025 – 5.943	0.00000350	3.334	3.209	3.458	0.993	a+
<i>Myctophum punctatum</i>	28	16 – 69	0.042 – 4.582	0.00000545	3.221	3.131	3.311	0.995	a+
<i>Notolychnus valdiviae</i>	27	15 – 22	0.019 – 0.123	0.00000181	3.484	2.342	3.859	0.410	i
<i>Notoscopelus bolini</i>	11	21 – 28	0.085 – 0.221	0.00000719	3.101	3.484	1.720	0.905	i
<i>Notoscopelus resplendens</i>	20	20 – 70	0.065 – 4.832	0.00000837	3.108	2.894	3.320	0.981	i
<i>Polyipnus polli</i>	34	16 – 43	0.072 – 2.313	0.00000227	3.723	3.463	3.982	0.964	a+
<i>Sternoptyx diaphana</i>	29	9 – 41	0.014 – 1.861	0.00001472	3.104	2.900	3.308	0.973	i
<i>Stomias boa boa</i>	20	53 – 153	0.330 – 8.212	0.00000158	3.042	2.781	3.302	0.971	i
<i>Sigmops elongatus</i>	35	45 – 151	0.165 – 7.879	0.00000075	3.205	3.021	3.387	0.975	a+
<i>Vinciguerrria nimbaria</i>	32	14 – 50	0.027 – 1.339	0.00000416	3.245	3.156	3.333	0.995	a+

**TABLE S2** Length-weight parameters for mesopelagic species (ordered alphabetically) from the equatorial and tropical Atlantic waters. *a*, intercept; *a*+, positive allometry; *a*-, negative allometry; *b*, allometry coefficient (slope); *WW*, eviscerate body wet weight; *i*, isometry; *n*, sample size; *SL*, standard length; *r*<sup>2</sup>, correlation of determination; 95% CL of *b*, confidence interval

Species	n	Range <i>SL</i> (mm)	Range <i>eWW</i> (g)	<i>a</i>	<i>b</i>	Inferior 95% CL of <i>b</i>	Superior 95% CL of <i>b</i>	<i>r</i> <sup>2</sup>	Growth model
<i>Argyropelecus sladeni</i>	27	15 – 43	0.053 – 0.910	0.00000844	3.208	2.681	3.735	0.863	i
<i>Benthoosema glaciale</i>	15	15 – 35	0.027 – 0.528	0.00000367	3.323	3.179	3.467	0.991	a+
<i>Benthoosema suborbitale</i>	29	16 – 33	0.060 – 0.478	0.00001402	2.993	2.584	3.401	0.943	i
<i>Bolinichthys photothorax</i>	6	21 – 51	0.116 – 1.520	0.00000994	3.032	2.611	3.452	0.990	i
<i>Ceratoscopelus warmingii</i>	47	16 – 64	0.033 – 3.368	0.00000364	3.303	3.214	3.391	0.992	a+
<i>Chauliodus danae</i>	43	25 – 212	0.059 – 22.652	0.00000266	2.957	2.855	3.059	0.988	i
<i>Cyclothone acclinidens</i>	22	31 – 46	0.033 – 0.274	0.00004776	2.183	1.738	2.635	0.774	a-
<i>Cyclothone livida</i>	32	23 – 37	0.039 – 0.146	0.00001491	2.538	2.032	3.045	0.777	i
<i>Cyclothone pallida</i>	24	25 – 54	0.047 – 0.380	0.00001526	2.540	2.135	2.944	0.885	a-
<i>Cyclothone pseudopallida</i>	19	23 – 35	0.027 – 0.083	0.00005324	1.986	1.178	2.794	0.613	a-
<i>Diaphus brachycephalus</i>	100	12 – 47	0.011 – 3.175	0.00000692	3.285	3.164	3.406	0.967	a+
<i>Diaphus dumerilii</i>	30	31 – 57	0.387 – 2.399	0.00001871	2.928	2.716	3.140	0.966	i
<i>Diaphus fragilis</i>	10	46 – 79	0.748 – 3.857	0.00000711	3.019	2.799	3.239	0.992	i
<i>Diaphus holti</i>	91	11 – 50	0.012 – 2.102	0.00000965	3.134	3.002	3.265	0.962	a+
<i>Diaphus metopoclampus</i>	10	19 – 40	0.064 – 0.802	0.00000330	3.349	2.979	3.718	0.982	i
<i>Diaphus mollis</i>	30	29 – 55	0.358 – 2.678	0.00001476	3.016	2.857	3.175	0.983	i
<i>Diaphus problematicus</i>	9	45 – 72	0.869 – 3.234	0.00001349	2.888	2.473	3.304	0.975	i
<i>Diaphus rafinesquii</i>	125	11 – 70	0.026 – 5.829	0.00002696	2.864	2.802	2.925	0.986	a-
<i>Diaphus vanhoeffeni</i>	17	25 – 36	0.248 – 0.584	0.00018969	2.238	1.740	2.735	0.860	a-
<i>Hygophum macrochir</i>	30	20 – 42	0.097 – 1.047	0.00001658	2.964	2.806	3.122	0.981	i
<i>Lampanyctus alatus</i>	142	16 – 57	0.040 – 1.720	0.00000418	3.214	3.121	3.306	0.971	a+
<i>Lampanyctus nobilis</i>	19	28 – 74	0.128 – 3.097	0.00000311	3.224	2.965	3.483	0.976	i
<i>Lampanyctus pusillus</i>	22	24 – 36	0.079 – 0.448	0.00000230	3.389	2.916	3.861	0.918	i
<i>Lepidophanes guentheri</i>	35	19 – 61	0.025 – 2.331	0.00000156	3.443	3.212	3.674	0.965	a+
<i>Lobianchia dofleini</i>	30	13 – 30	0.030 – 0.464	0.00000849	3.184	2.855	3.512	0.934	i
<i>Myctophum affine</i>	34	14 – 47	0.021 – 1.491	0.00000254	3.440	3.291	3.589	0.986	a+
<i>Myctophum nitidulum</i>	25	15 – 74	0.021 – 5.137	0.00000243	3.400	3.281	3.518	0.994	a+
<i>Myctophum punctatum</i>	28	16 – 69	0.036 – 3.863	0.00000438	3.232	3.137	3.326	0.995	a+
<i>Notolychnus valdiviae</i>	27	15 – 22	0.019 – 0.123	0.00000278	3.303	2.498	3.759	0.424	i
<i>Notoscopelus bolini</i>	11	21 – 28	0.077 – 0.195	0.00000594	3.129	3.302	1.587	0.933	i
<i>Notoscopelus resplendens</i>	20	20 – 70	0.060 – 4.427	0.00000756	3.114	2.908	3.320	0.983	i
<i>Polyipnus polli</i>	34	16 – 43	0.060 – 2.070	0.00000207	3.719	3.453	3.985	0.962	a+
<i>Sternoptyx diaphana</i>	29	9 – 41	0.009 – 1.604	0.00000729	3.236	3.038	3.434	0.977	a+
<i>Stomias boa boa</i>	20	53 – 153	0.312 – 7.306	0.00000159	3.013	2.739	3.292	0.967	i
<i>Sigmops elongatus</i>	35	45 – 151	0.147 – 7.149	0.00000064	3.220	3.064	3.376	0.982	a+
<i>Vinciguerrria nimbaria</i>	32	14 – 50	0.023 – 0.964	0.00000594	3.073	2.989	3.158	0.995	i